

On the scaling properties of $2d$ randomly stirred Navier–Stokes equation

Andrea Mazzino*

*Department of Physics, University of Genova, and INFN, CNISM,
Genova Section, Via Dodecaneso 33, I-16146 Genova, Italy,*

Paolo Muratore-Ginanneschi†

Department of Mathematics and Statistics, University of Helsinki, P.O. Box 4, 00014 Helsinki, Finland,

Stefano Musacchio‡

*Department of Physics of Complex Systems Weizmann Institute of Science Rehovot 76100, Israel
(Dated: February 8, 2008)*

We inquire the scaling properties of the $2d$ Navier-Stokes equation sustained by a forcing field with Gaussian statistics, white-noise in time and with power-law correlation in momentum space of degree $2 - 2\varepsilon$. This is at variance with the setting usually assumed to derive Kraichnan’s classical theory. We contrast accurate numerical experiments with the different predictions provided for the small ε regime by Kraichnan’s double cascade theory and by renormalization group (RG) analysis. We give clear evidence that for all ε Kraichnan’s theory is consistent with the observed phenomenology. Our results call for a revision in the RG analysis of $(2d)$ fully developed turbulence.

PACS numbers: PACS number(s): 47.27.Te, 47.27.–i, 05.10.Cc

In two dimensions, the joint conservation of energy and enstrophy has relevant consequences for the Navier–Stokes equation. In $2d$ it is possible to prove that in the deterministic case the solution of the Cauchy problem exists and is unique [20] and, very recently, that in the stochastic case [8, 9, 10, 12, 19, 23] the solution is a Markov process exponentially mixing in time and ergodic with a unique invariant (steady state) measure even when the forcing acts only on two Fourier modes [22]. At large Reynolds number, the $2d$ scaling properties also differ from the $3d$ -case. A long standing hypothesis [25], very recently corroborated by numerical experiments [6, 7], surmises the existence of a conformal invariance in $2d$. A phenomenological theory due to Kraichnan [18] and Batchelor [2] predicts the presence of a double cascade mechanism governing the transfer of energy and enstrophy in the limit of infinite inertial range. Accordingly, an *inverse energy cascade* with spectrum characterized by a scaling exponent $d_\varepsilon = -5/3$ appears for values of the wave-number k smaller than the typical scale k_F of the forcing scale. For wave-numbers larger than k_F a *direct enstrophy cascade* should occur. The corresponding energy spectrum scales as $d_\varepsilon = -3 + \dots$ where the dots here stand for possible logarithmic corrections. As emphasized in [3, 5] Kraichnan’s theory is encoded in three hypothesis, (i) velocity correlations are smooth at finite viscosity and exist in the inviscid limit even at coinciding points, (ii) Galilean invariant functions and in particular structure functions reach a steady state and (iii) no dissipative anomalies occur for the energy cascade. Under these hypotheses, if the forcing field is homogeneous and isotropic Gaussian and time δ -correlated it is possible to derive asymptotic expressions of the three point structure functions of the velocity field consistent with Kraich-

nan’s predictions [3, 5, 21]. Very strong laboratory (see [17, 26] and reference therein) and numerical evidences (see e.g. [11] and references therein) support Kraichnan’s theory. However, a first principle derivation of the statistical properties of two dimensional turbulence is still missing. An attempt in this direction has recently been undertaken [14, 15] (see also [24] and [16]) by applying a renormalization group improved perturbation theory (RG) ([27] and [1] for review of applications to fluid turbulence) to the randomly stirred Navier–Stokes equation with power law forcing. However how it will be discussed in details in the sequel, RG analysis leads to a scenario not a priori consistent with Kraichnan’s theory. The goal of the present work is to shed light on this issue.

RG starting point is the randomly stirred Navier–Stokes equation

$$(\partial_t + v \cdot \partial)v^\alpha - \nu_0 \partial^2 v^\alpha = -\partial^\alpha P + f^\alpha - \xi_0 v^\alpha \quad (1)$$

$$\partial \cdot v = 0$$

where $\alpha = 1, 2$, P is the pressure enforcing incompressibility and ξ_0 is an Eckman type coupling providing for large scale dissipation. The forcing f is a Gaussian field with zero average and correlation

$$\prec f^\alpha(\mathbf{x}, t) f^\beta(\mathbf{y}, s) \succ = \delta(t - s) F^{\alpha\beta}(\mathbf{x} - \mathbf{y}) \quad (2)$$

$$F^{\alpha\beta}(\mathbf{x}) := \int \frac{d^d p}{(2\pi)^d} e^{i\mathbf{p} \cdot \mathbf{x}} \tilde{F}(p) T^{\alpha\beta}(\hat{\mathbf{p}}) \quad (3)$$

$$\tilde{F}(p) := \frac{F_0 \chi\left(\frac{p}{m}, \frac{p}{M}\right)}{p^{d-4+2\varepsilon}} \quad (4)$$

F_0 is a constant specifying the amplitude of forcing fluctuations and $T^{\alpha\beta}(p)$ is the transversal projector. The function χ in (4) is slowly varying for $m \ll p \ll M$ and set infra-red m and ultra-violet cut-offs M scales for the

forcing. Its detailed shape does not affect the scaling predictions of the RG. Irrespectively of the spatial dimension d , the cumulative spectrum $F_\alpha^\alpha(0)$ of the forcing (Einstein convection for repeated vector indexes) diverges for $0 \leq \varepsilon < 2$ as the ultra-violet cut-off M tends to infinity hence providing for stirring at small spatial scales. For $\varepsilon > 2$, $F_\alpha^\alpha(0)$ is dominated by small wave-numbers and thus describes infra-red stirring. At $\varepsilon = 0$, the scaling dimensions of the material derivative, dissipation and forcing in (1) coincide for a scaling dimension of the velocity field $d_v = 1$ in momentum units. Thus $\varepsilon = 0$ provides a marginal limit around which scaling dimensions can be perturbatively determined with the help of ultra-violet RG. The main result [14, 15, 16] is the existence of a non-Gaussian infra-red stable fixed point of the RG flow yielding for the energy spectrum the prediction

$$\mathcal{E}(k) = \varepsilon^{1/3} F_0^{2/3} k^{1-\frac{4\varepsilon}{3}} R\left(\varepsilon, \frac{m}{k}, \frac{k_b}{k}\right) \quad (5)$$

The adimensional function R depends upon infra-red scales m and $k_b = (\varepsilon \xi_0^3 / F_0)^{1/(6-2\varepsilon)}$ and admits a regular expansion in powers of ε at the RG fixed point [14]. The resummation leading to $k^{1-\frac{4\varepsilon}{3}}$ in (5) is derived from the solution of the Callan–Symanzik equation [27] which in the present case requires scaling at finite ε to stem from the balance of the two terms in the material derivative with the forcing i.e. from the requirement of Galilean invariance alone. According to [1, 14, 15] composite operators at small ε does not induce any self-similarity breaking by the infra-red scales m and k_b . The conclusion is that the spectrum should scale for small ε with exponent $d_\mathcal{E} = 1 - 4\varepsilon/3$ as in the 3d-case [1]. Such conclusion seems at variance with Kraichnan’s theory which instead suggests the occurrence of an inverse cascade at small ε . Namely under the assumptions (i),(ii),(iii) [3, 4] the energy balance equation

$$\begin{aligned} (\partial_t + \xi_0 - \nu_0 \partial^2) \prec v^\alpha(\mathbf{x}, t) v_\alpha(0, t) \succ \\ - \frac{1}{2} \partial_\mu \prec ||\delta v||^2(\mathbf{x}, t) \delta v^\mu(\mathbf{x}, t) \succ = F(\mathbf{x}) \end{aligned} \quad (6)$$

with $\delta \mathbf{v}^\alpha(\mathbf{x}, t) = v^\alpha(\mathbf{x}, t) - v^\alpha(0, t)$ yields for $\xi_0 = 0$, $\varepsilon < 2$ and $r^2 := ||x||^2$

$$S_3(r) \simeq c_1 F_0 M^{4-2\varepsilon} r + \frac{c_2 F_0}{r^{3-2\varepsilon}}, \quad m^{-1} \gg r \gg M^{-1} \quad (7)$$

S_3 denotes the three point velocity longitudinal structure function and c_i $i = 1, 2$ two adimensional coefficients irrelevant for the present argument. Power counting based on (7) then predicts a $d_\mathcal{E} = -5/3$ inverse cascade spectrum at small ε with *at most sub-leading* corrections consistent with the RG prediction.

The currently available numerical resources permit to contrast the two apparently discordant predictions (5) and (7) with the actual Navier–Stokes phenomenology.

To attain this goal we integrated the Navier–Stokes equation (1) for the vorticity field ($\omega = \epsilon_{\alpha\beta} \partial^\beta v^\alpha$) with a standard, fully-dealiased pseudospectral method in a doubly periodic square domain of resolution 1024^2 . Time evo-




FIG. 1: Energy spectrum for $\varepsilon = 0.5$ and standard molecular dissipation. An inertial range with inverse energy cascade sets in for sufficiently small viscosity $\nu = O(10^{-4})$ (solid line). At larger viscosity the inertial range is suppressed and dissipative spectrum $\mathcal{E} \sim k^{1-2\varepsilon}$ is observed (dashed line).

lution was computed by means of a standard second-order Runge–Kutta scheme. We repeated our numerical experiments for different values of the hyperviscosity $(-1)^{p+1} \nu_0 \partial^{2p} v$ including $p = 1$, standard viscosity as in (1), and $p = 4, 6$ obtaining qualitatively identical results. The outcomes evince that for *all* $0 \leq \varepsilon \leq 2$ and sufficiently small viscosity, an inverse energy cascade with exponent consistent with $d_\mathcal{E} = -5/3$ is observed (see Fig. 1). If the viscosity increases, finite resolution effects prevent the observation of an inertial range as the Kolmogorov scale becomes of the order of the infra-red dissipation scale. In such a case a dissipative spectrum appears with scaling exponent consistent with the prediction $d_\mathcal{E} = 1 - 2\varepsilon$ dictated by the balance between forcing and dissipation (see again Fig. 1). In agreement with (7), local balance scaling becomes dominant in the range $2 < \varepsilon < 3$ (see Fig. 2). There, energy spectra scale with an exponent compatible with $d_\mathcal{E} = 1 - 4\varepsilon/3$. Finally, for $\varepsilon > 3$ a direct enstrophy cascade invades the whole inertial range. The phenomenology just described is confirmed by the inspection of the energy and enstrophy fluxes in the inertial range (see Fig. 3). They are respectively defined by $\Pi_E(k) \propto \int_{k_o}^k \frac{d^2 q}{(2\pi)^2} \Re \prec \mathcal{F}\{v\}^\alpha (-q) \mathcal{F}\{(v \cdot \partial v_\alpha)\}(q) \succ$ and $\Pi_Z(k) \propto \int_{k_o}^k \frac{d^2 q}{(2\pi)^2} \Re \prec \mathcal{F}\{\omega\} (-q) \mathcal{F}\{(v \cdot \partial \omega)\}(q) \succ$ where $\mathcal{F}$ denotes the Fourier transform and k_o is chosen such that $\Pi_E(k)$ and $\Pi_Z(k)$ are positive quantities in the inertial range. As spectra and structure functions, fluxes highlight the existence versus ε of three distinct scaling regimes the origin of which can be understood by contrasting (6) with the energy and enstrophy inputs




FIG. 2: Third order structure function of longitudinal velocity increments $S_3(r)$ for $\epsilon = 1$ (squares), $\epsilon = 2.5$ (circles), $\epsilon = 4$ (triangles). When $\epsilon < 2$ the presence of an inverse energy cascade with constant flux results in the scaling $S_3(r) \sim r$ (solid line). When $\epsilon > 3$ the presence of a direct enstrophy cascade with constant flux results in the scaling $S_3(r) \sim r^3$ (dashed line). For $2 < \epsilon < 3$ the scaling is compatible with the RG argument (dash-dotted line)

$I_E(k) \propto \int_{k_0}^k dq q \tilde{F}_v(q)$ and $I_Z(k) \propto \int_{k_0}^k dq q \tilde{F}_v(q)$. For $0 \leq \epsilon < 2$ both I_E and I_Z are peaked in the ultra-violet. Since stirring occurs mainly at small spatial scales an inverse energy cascade sets in the whole inertial range. In the range $2 < \epsilon < 3$ the energy input becomes infra-red divergent whilst I_Z is still ultra-violet divergent in the absence of cut-offs. In this situation, the steady state is attained when the energy and enstrophy fluxes balance scale by scale the corresponding inputs (see Fig. 3). In this ϵ -range the energy spectrum scales in agreement with the exponent which can be extrapolated but no longer fully justified using the RG. Finally, for $\epsilon > 3$ both the energy and enstrophy inputs become peaked in the infra-red. In such a case [3, 4] the right hand sides of Eq. (6) and of the analogous equation for the vorticity correlation admit a regular Taylor expansion for $mx \ll 1$. The hypotheses (i), (ii), (iii) thus recover $S_3(x) \propto r^3$ i.e. Kraichnan's scaling for the *direct enstrophy cascade* in agreement with our numerical observations.

As far as intermittency is concerned, our numerics support normal scaling of velocity structure functions (Fig. 4). Higher order vorticity structure function are compatible with a weakly anomalous scaling (Fig. 5). These results provide an indirect positive test for the occurrence of a vorticity dissipative anomaly [4].

In conclusion, our numerics fully support the validity of Kraichnan's theory for all values of the Hölder exponent of power law forcing in $2d$. RG scaling prediction is not observed in the range where it was supposed to appear. It seems to us that this is not trivially a consequence of low Reynolds number ($O(\epsilon^{1/2})$ for $F_0 = O(\epsilon)$ see discussion in section 9.6.4 of [13]) entailed by perturbative expansion. Instead scaling seems to be related




FIG. 3: The energy flux Π_E (symbols) and the energy input $I_E(k)$ (thin line) for different values of ϵ . (a) $\epsilon = 1$; (b) $\epsilon = 2.5$; (c) $\epsilon = 4$. The energy input is defined as $I_E(k) \propto \int_0^k dq q \tilde{F}_v(q)$ for $\epsilon < 2$ and as $I_E(k) \propto \int_k^\infty dq q \tilde{F}_v(q)$ for $\epsilon > 2$. In (b) the enstrophy input, $I_Z(k) \propto \int_0^k dq q^3 \tilde{F}(q)$ with and the enstrophy flux $\Pi_Z(k)$ are also plotted in order to show their balance at each inertial range scale.

to the convolution of the response field with the forcing kernel in the Schwinger-Dyson equation [1, 27] giving rise to a non-local scaling field of dimension $d_O = -1/3$ independently of ϵ . This point however deserves more theoretical inquiry.

We are pleased to thank J. Honkonen, A. Kupiainen and P. Olla for numerous friendly and enlightening discussions. Numerical simulations have been performed on the “Turbofarm” cluster at the INFN computing center in Torino. A.M. was supported by COFIN 2006 N.2005027808 and by CINFAl consortium, P.M.-G. by centre of excellence “Geometric Analysis and Mathematical Physics” of the Academy of Finland and by FP5 EU network contract HPRN-CT-2002-00300.




FIG. 4: Structure function of longitudinal velocity increments $S_4(r)$ (squares) and $S_6(r)$ (circles) vs. $S_2(r)$. The lines represent the non-intermittent scalings $S_4(r) \sim S_2(r)^2$ (solid line) and $S_6(r) \sim S_2(r)^3$ (dashed line). (a) $\epsilon = 1$; (b) $\epsilon = 2.5$; (c) $\epsilon = 4$




FIG. 5: Structure function of vorticity increments $S_n^{(\omega)}(r)$ for $n = 1, 2, 3, 4$ (square, circles, triangles, poligons). In the inset we show the scaling exponents ζ_n . Here $\epsilon = 4$.

- [4] D. Bernard, SPhT-00-039 and [chao-dyn/9904034](#).
- [5] D. Bernard, SPhT-00-093 and [cond-mat/0007106](#).
- [6] D. Bernard, G. Boffetta, A. Celani and G. Falkovich, *Nature Physics* **2** 124 (2006)
- [7] D. Bernard, G. Boffetta, A. Celani and G. Falkovich, [nlin.CD/0609069](#)
- [8] J. Bricmont, A. Kupiainen and R. Lefevre, *J. Stat. Phys.*, **100** 743 (2000) and [math-ph/9912018](#).
- [9] J. Bricmont, A. Kupiainen and R. Lefevre, *Comm. Math. Phys.* **224** 65 (2001) and [www.ma.utexas.edu/mp_arc/](#)
- [10] J. Bricmont, A. Kupiainen and R. Lefevre, *Comm. Math. Phys.* **230** 87 (2002) and [math-ph/0010028](#).
- [11] G. Boffetta, submitted to *Phys. Fluids* (2006).
- [12] W. E, J. C. Mattingly, and Y. G. Sinai. *Comm. Math. Phys.* **224** 83 (2001).
- [13] U. Frisch, *Turbulence*, Cambridge Univ. Press, Cambridge, (1995).
- [14] J. Honkonen, *Phys. Rev. E* **58** 4532 (1998).
- [15] J. Honkonen, Yu.S. Kabrits and M.V. Kompaniets, [nlin/0201025](#).
- [16] L. Ts. Adzhemyan, J. Honkonen, M. V. Kompaniets and A. N. Vasilév, *Phys. Rev. E* **71**, 036305 (2005) and [nlin.CD/0407067](#).
- [17] H. Kellay and W.I. Goldburg, *Rep. Prog. Phys.* **65** 845 (2002).
- [18] R. Kraichnan, *Phys. Fluids* **10** (1967) 1417.
- [19] S. Kuksin and A. Shirikyan. *Comm. Math. Phys.* **213** 291 (2000).
- [20] O. A. Ladyzhenskaya *The Mathematical Theory of Viscous Incompressible Flow*, Gordon and Breach, New York, 1969.
- [21] E. Lindborg, *Phys. Fluids* **11** 510 (1999).
- [22] M. Hairer and J.C. Mattingly, *Annals of Mathematics*, **164** (2006) 993 and [math.PR/0406087](#).
- [23] J.C. Mattingly, *Comm. Math. Phys.* **230** 421 (2002).
- [24] P. Olla, *Phys. Rev. Lett.* **67** 2465 (1991).
- [25] A. Polyakov, *Nucl.Phys. B* **396** 367 (1993) and [hep-th/9212145](#)
- [26] P. Tabeling, *Physics Reports* **362** 1 (2002)
- [27] J. Zinn-Justin, *Quantum field theory and critical phenomena*, 4th ed. Clarendon Press, Oxford (2002).

* Electronic address: mazzino@fisica.unige.it

† Electronic address: paolo.muratore-ginanneschi@helsinki.fi

‡ Electronic address: stefano.musacchio@gmail.com

- [1] L.Ts. Adzhemyan, A.V. Antonov and A.N. Vasilév *The field theoretic renormalization group in fully developed turbulence* Gordon and Breach, Amsterdam (1999).
- [2] G.K. Batchelor, *Phys. Fluids*, **Supp II 12** 233 (1969).
- [3] D. Bernard, *Phys. Rev E* **60** 6184 (1999) and [chao-dyn/9902010](#).